Bull. Southern California Acad. Sci.
96(3), 1997, pp. 117–121
© Southern California Academy of Sciences, 1997

Shortfin Mako, *Isurus oxyrinchus*, Impaled by Blue Marlin, *Makaira nigricans* (Teleostei: Istiophoridae)

Harry L. Fierstine,[1] Gregor M. Cailliet,[2] and Julie A. Neer[2]

[1]*Biological Sciences Department, California Polytechnic State University San Luis Obispo, CA 93407*
[2]*Moss Landing Marine Laboratories, Moss Landing, CA 95039-0450*

Although billfish are known to drive their rostra into pelagic sharks, most accounts attribute impalement to the swordfish (*Xiphias gladius*) and not to a member of the family Istiophoridae (Gudger 1940; Smith 1961; Starck 1960). In the only record of a shark having been stabbed by an istiophorid billfish, Cliff et al. (1990) reported that a female shortfin mako (*Isurus oxyrinchus*) was caught off South Africa with the broken rostrum of a sailfish (*Istiophorus platypterus*) embedded in its left orbit. Few other details were given. We present an account of a shortfin mako caught off Baja California, Mexico that was discovered during fish processing operations to have an istiophorid rostrum embedded in its vertebral column (Fig. 1). We briefly discuss the effects of impalement on both shark and billfish.

According to David Arpia (American Sea Food Co., Inc., San Diego, CA, April 3, 1995), "the shark, a male, was caught by longline about 72 km west of Enseñada, Baja California, Mexico, and weighed approximately 28.3 kg dressed. Length was not recorded. It appeared healthy at time of capture with only a slight blemish in its skin where the rostrum entered the body. Except for a capsule of scar tissue surrounding the bone fragment, the flesh appeared normal, and was processed for sale as usual. The broken (posterior) end of the bill fragment was located about two inches (5 cm) below the skin surface." The bill entered the left side of the shark at the level of the second dorsal fin, completely penetrating a vertebral centrum without entering muscle tissue of the right side. A section of the vertebral column with the embedded bill (SIO 94-4) is housed in the Marine Vertebrates Collection, Scripps Institution of Oceanography, La Jolla, CA 92093.

We follow the methodology and terminology of Fierstine and Voigt (1996) and Fierstine and Crimmen (1996) in identifying rostral fragments. Because the bill in this study was a distal segment, measurements and ratios were compared with values from Table 1 of Fierstine and Voigt (1996) at one-fourth bill length (0.25L, or one-fourth the distance between the tip and the orbital margin of the lateral ethmoid).

The bill fragment (Table 1; Fig. 1) is from the distal end of the rostrum and is worn slightly at its tip. We estimate that it was broken slightly anterior to 0.25 L, with paired nutrient canals exposed on the posterior surface. The rostrum is from *Makaira nigricans* (blue marlin) because of the following combination of features: (1) placement of the nutrient canals (DD/D) with respect to depth of the rostrum at 0.25L (=0.65, a value outside the observed range of all istiophorids, but very close to *M. nigricans* and *Tetrapturus angustirostris*); (2) the absence of denticles on the dorsal surface (eliminates all istiophorids inhabiting the Indo-Pacific Ocean except *M. nigricans* and *T. angustirostris*); (3) no evidence of a prenasal in the fragment (eliminates *T. angustirostris*).



Fig. 1. Vertebral segment at level of second dorsal fin of *Isurus oxyrinchus* (SIO 94-4) with embedded rostral fragment of *Makaira nigricans*.

Rostrum morphology suggests the blue marlin was similar in size to Los Angeles County Museum specimen LACM 25491, i.e. slightly less than 44 kg and 173 cm lower jaw to fork length. Blue marlin of this small size would be an immature fish (Hopper 1986).

To obtain an estimate of total length (TL) of the mako shark, we used several morphometric conversion equations developed from second dorsal fin (Garrick 1967; Gubanov 1974; Applegate 1977) and centrum diameter (Cailliet et al. 1983a) measurements in the literature, and from one female shortfin mako with a similar-sized second dorsal fin collected off southern California. Both methods of estimating length resulted in ranges between 200–230 cm TL. We follow the methodology of Cailliet et al. (1983a, b) in estimating age of the shark. Two vertebrae were removed from the vertebral column directly below the second dorsal fin, one immediately anterior and another posterior to insertion of the bill. Age was estimated by counting the number of opaque bands observed in the x-

Table 1. Measurements and ratios of a rostral fragment of *Makaira nigricans* found impaled in *Isurus oxyrinchus* (SIO 94-4).

Measurements (mm)					Ratios		
Length of fragment	Width (W) @ 0.25L	Depth (D) @ 0.25L	Height of nutrient canals (H) @ 0.25L	Distance of nutrient canal from dorsum(DD) @ 0.25L	D/W	H/D	DD/D
98	14.4	9.3	1.0	6.0	0.65	0.11	0.65

rays and by counting ridges along the anterior surface of a whole vertebra. Two different age estimates were produced, one using the assumption of one band pair per year (Cailliet et al. 1983a) and the other using two bands pairs per year (Pratt and Casey 1983a, b).

Both vertebral analysis techniques produced estimates of 7 or 8 band pairs. Thus, the age of this mako was either 3–4 years (Pratt and Casey 1983a, b) or 7–8 years (Cailliet et al. 1983a). A male shortfin mako of this size (200–230 cm TL) would be close to mature (Cailliet et al. 1983a), regardless of age. Female shortfin makos become mature at a larger size, approximately 280 cm TL (Stevens 1983).

There is no evidence to determine if the place of capture of the shortfin mako was also the site of its encounter with the blue marlin. The presence of both *I. oxyrinchus* and *M. nigricans* is more commonly found off Baja California Sur than Baja California. Both species favor a similar epipelagic, oceanic environment, have a worldwide distribution (Compagno 1984; Eschmeyer et al. 1983), and are capable of long distance migrations (Nettles et al. 1994; National Marine Fisheries Service, unpublished).

When the blue marlin broke away from the shark, we assume fracture occurred at the skin surface (the thinnest region of the exposed rostrum) and not within the muscle. Friction between the denticles and the cartilage and muscle tissue probably kept the rostrum from being withdrawn. As both fish struggled to separate, the skin surface functioned as a fulcrum about which the bill fractured in a dorso-ventral plane. Since the rostrum was found 5 cm below the skin surface, the shark grew in caudal diameter (and presumably length) after it had been speared. When compared with vertebrae from an unimpaled shortfin mako of similar size, band patterns did not indicate an alteration of growth after stabbing.

Shortfin makos are opportunistic predators often feeding on scombrids (Compagno 1984) and occasionally on istiophorids, and generally eat prey between 23–35% of their length (Cliff et al. 1990). The sailfish responsible for spearing the mako shark off South Africa (Cliff et al. 1990) was well within the size range of prey consumed by shortfin mako, thus the spearing probably was a defensive response to a predatory act. In contrast, the incident we present here may not have occurred during feeding because the blue marlin probably exceeded the preferred prey size of the shark. We do not know the exact length of either fish at time of impalement but the blue marlin (TL = ~173 cm lower jaw to fork + tip of lower jaw to tip of bill + fork to tip of caudal fin) was probably as long as or longer than the shortfin mako (TL < 200–230 cm).

Most effects of physical encounters between sharks and billfish are unknown. The shortfin mako that was captured off South Africa was impaled in its left orbit and was underweight for its length, and Cliff et al. (1990) speculated that the shark's vision had been impaired enough to affect foraging success. In the case presented here, the shark apparently remained healthy because no vital organ was pierced. In contrast, we know little of the effect on billfish after a rostral fracture. It seems reasonable that loss of a section of the bill would make a billfish more vulnerable to predation, but evidence is lacking. We do know some fish recover from their injuries because there are several records of apparently healthy billfishes with damaged, malformed, or missing rostra (Frazier et al. 1994).

Acknowledgments

We thank R. N. Lea and C. R. Robins for reviewing an earlier version of the manuscript, and R. H. Rosenblatt for bringing the speared shortfin mako to our attention. G. Cliff graciously provided references and allowed us to examine the sailfish rostrum he removed from the shortfin mako.

Literature Cited

American Fisheries Society. 1991. Common and scientific names of fishes from the United States and Canada. 5th ed. Amer. Fish. Soc. Spec. Pub., 20:1–183.

Applegate, S. P. 1977. A new record-size bonito shark, *Isurus oxyrinchus* Rafinesque, from southern California. Calif. Fish and Game, 63:126–129.

Cailliet, G. M., L. K. Martin, J. T. Harvey, D. Kusher, and B. A. Welden. 1983a. Preliminary studies on the age and growth of blue, *Prionace glauca*, common thresher, *Alopias vulpinus*, and shortfin mako, *Isurus oxyrinchus*, sharks from California waters. Pp. 179–188 *in* Proc. of the international workshop on age determination of oceanic pelagic fishes: Tunas, billfishes, and sharks (E. D. Prince and L. M. Pulos, eds.). U.S. Dept. Commer., NOAA Tech. Rep. NMFS 8.

———, L. K. Martin, D. Kusher, P. Wolf, and B. A. Welden. 1983b. Techniques for enhancing vertebral bands in age estimation of California elasmobranchs. Pp. 157–165 *in* Proc. of the international workshop on age determination of oceanic pelagic fishes: Tunas, billfishes, and sharks (E. D. Prince and L. M. Pulos, eds.). U.S. Dept. Commer., NOAA Tech. Rep. NMFS 8.

Cliff, G., S. F. J. Dudley, and B. Davis. 1990. Sharks caught in the protective gill nets off Natal, South Africa. 3. The shortfin mako shark, *Isurus oxyrinchus* (Rafinesque). South African J. of Marine Sci., 9:115–126.

Compagno, L. J. V. 1984. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. FAO Fisheries Synopsis no. 125., 4(1):i–viii+1–249.

Eschmeyer, W. N., E. S. Herald, and H. Hammann. 1983. A field guide to Pacific Coast fishes of North America from the Gulf of Alaska to Baja California. Houghton Mifflin Co., xiv+336 pp.

Fierstine, H. L., and N. L. Voigt. 1996. Use of rostral characters for identifying adult billfishes (Teleostei: Perciformes: Istiophoridae and Xiphiidae). Copeia, 1996:148–161.

———, and O. Crimmen. 1996. Two erroneous, commonly cited examples of "swordfish" piercing wooden ships. Copeia, 1996:472–475.

Frazier, J. G., H. L. Fierstine, S. C. Beavers, F. Achaval, H. Suganuma, R. L. Pitman, Y. Yamaguchi, and C. M. Prigioni. 1994. Impalement of marine turtles (Reptilia, Chelonia: Cheloniidae and Dermochelyidae) by billfishes (Osteichthyes, Perciformes: Istiophoridae and Xiphiidae). Env. Biol. Fishes, 39:85–96.

Garrick, J. A. F. 1967. Revision of sharks of genus *Isurus* with description of a new species (Galeoidea, Lamnidae). Proc. U.S. Natl. Mus., 118:663–690.

Gubanov, Y. P. 1974. The capture of a giant specimen of the mako shark (*Isurus glaucus*) in the Indian Ocean. J. Ichthyol., 14:589–591.

Gudger, E. W. 1940. The alleged pugnacity of the swordfish and the spearfishes as shown by their attacks on vessels. Mem. Royal Asiatic Soc. Bengal, 12:215–315.

Hopper, C. N. 1986. The majestic marlin solves mysteries of the sea. Pp. 58–61, 63, 65–66, 68–70 *in* World Record Game Fishes 1986,. International Game Fish Association, Fort Lauderdale, Florida.

Leviton, A. E., R. H. Gibbs Jr., E. Heal, and C. D. Dawson. 1985. Standards in herpetology and ichthyology: Part I. Standard symbolic codes for institutional resource collections in herpetology and ichthyology. Copeia, 1985:802–832.

Nakamura, I. 1985. An annotated and illustrated catalogue of marlins, sailfishes, spearfishes and swordfishes known to date. FAO Fisheries Synopsis no. 125., 5:i–iv+1–65.

National Marine Fisheries Service. unpublished. The Shark Tagger 1992 Summary, Newsletter of the Cooperative Shark Tagging Program, Northeast Fisheries Science Center, Narragansett, RI 02882.

Nettles, C. I., R. E. Bayley, C. D. Jones, and M. T. Judge. 1994. Cooperative game fish tagging program annual newsletter: 1992. Pp. 1–23 *in* M. I. Farber and E. D. Prince (eds). U.S. Dept. Commerc., NOAA Tech. Mem. NMFS-SEFSC-36.

Pratt, H. L., Jr., and J. G. Casey. 1983a. Age and growth of the shortfin mako, *Isurus oxyrinchus*. Summary paper, pp. 175–177 *in* Proc. of the international workshop on age determination of oceanic pelagic fishes: Tunas, billfishes, and sharks (E. D. Prince and L. M. Pulos, eds.). U.S. Dept. Commerc., NOAA Tech. Rept. NMFS 8.

———. 1983b. Age and growth of the shortfin mako, *Isurus oxyrinchus*, using four methods. Canadian J. Fish. Aquat. Sci., 40:1944–1957.

Smith, J. L. B. 1961. Battle of sea-giants. Field and Tide, 3(9):6–7.

Starck, W. A. 1960. Spear of swordfish, *Xiphias gladius* Linnaeus, embedded in a silk shark, *Eulamia floridana* (Schroeder and Springer). Quart. J. Florida Acad. Sci., 23(2):165–166.

Stevens, J. D. 1983. Observations on reproduction in the shortfin mako, *Isurus oxyrinchus*. Copeia, 1983:126–130.

Accepted for publication 12 June 1997.